

**From Mutation Identification to Therapy: Discovery and Origins of the First Approved Gene Therapy in the Western World.**

John J. P. Kastelein<sup>1</sup>, Colin J. D. Ross<sup>2,3,4</sup>, and Michael R. Hayden<sup>3,4</sup>,

<sup>1</sup> Department of Vascular Medicine, Academic Medical Centre, University of Amsterdam, Amsterdam, The Netherlands

<sup>2</sup> Department of Pediatrics, University of British Columbia, Vancouver, BC, Canada

<sup>3</sup> Centre for Molecular Medicine and Therapeutics, Child and Family Research Institute, University of British Columbia, Vancouver, BC, Canada

<sup>4</sup> Department of Medical Genetics, University of British Columbia, Vancouver, BC, Canada

Please address correspondence to M.R.H. or J.J.P.K.

Dr. Michael Hayden, Centre for Molecular Medicine and Therapeutics, 950 West 28<sup>th</sup> Avenue, Vancouver, B.C., Canada V5Z-4H4, Email: mrh@cmmr.ubc.ca

Dr. John Kastelein, Department of Vascular Medicine, Academic Medical Centre, University of Amsterdam, the Netherlands, Meibergdreef 9, 1105 AZ Amsterdam. Email: J.J.Kastelein@amc.uva.nl

## The Road to Glybera: Discovery and Origins of the First Gene Therapy

On November 2, 2012, Glybera® (alipogene tipovarvec) was the first human gene therapy to receive long awaited market approval in the Western world. This important milestone is expected to open the door to additional gene therapies for the treatment of many diseases in the future. The development of this new gene therapy involved the dedicated contributions from many people from around the world.

The story of Glybera® began 26 years ago, in September 1986, in the laboratory of Dr. Michael Hayden at the University of British Columbia in Vancouver, B.C., Canada. Dr. Kastelein, a Dutch physician joined the lab and began participating in the Lipid Research Clinic with Dr. Hayden to diagnose and treat metabolic lipid disorders, while at the same time learning new laboratory techniques, such as DNA extractions and Southern blotting. One of the first patients, a young Canadian adolescent of 19 years, presented with severe hypertriglyceridemia, cutaneous xanthomas, chronic abdominal pain, and recurrent pancreatitis (Fig 1). The symptoms were consistent with a rare genetic disease, Lipoprotein Lipase Deficiency. Comparing the DNA sequence of the patient's LPL gene to the recently published LPL gene sequence (Wion et al., 1987), Drs. Hayden and Kastelein, as well as Drs. Sylvie Langlois, Samir Deeb, and John Brunzell, identified the first mutations in the LPL gene that caused LPL deficiency (Langlois et al., 1989) (Fig. 2). Drs. Hayden and Kastelein recall reviewing the results of the first crucial Southern blot, showing both a deletion and insertion on the paternal and maternal alleles in the patient's DNA, respectively, and recognizing that this gene defect was the cause of the disease.

Conventional lipid-lowering medications were ineffective for the treatment for LPL deficiency, and attempts to manage patient symptoms had focused on severe restrictions of dietary fat(Brunzell et al., 2001). However, the disease symptoms often persisted for LPL deficient patients on strict restrictions of dietary fat, leaving patients in severe pain with recurrent hospitalizations, and in some cases fatal pancreatitis (Rip et al., 2003; 2005; Gaudet et al., 2012b). A decision was made to do more for these patients – to embark on a journey that would improve the lives of patients that suffered from this devastating disease. Dr. Hayden began to envisage new therapeutic options for LPL deficient patients that were based on the LPL gene itself.

Drs. Hayden and Kastelein began to extensively characterize patients with novel mutations in the LPL gene from Canada, Holland and around the world (Devlin et al., 1990; Henderson et al., 1990; Monsalve et al., 1990; Bruin et al., 1992; Henderson et al., 1992; Ma et al., 1992; Bijvoet et al., 1994; Bruin et al., 1994; Jukema et al., 1996; Pimstone et al., 1996; Bijvoet et al., 1996a; 1996b; Groenemeijer et al., 1997b; Foubert et al., 1997a; Kastelein et al., 1998; Wittekoek et al., 1998; Henderson et al., 1998a; 1998b). In regions of the province of Quebec, Canada, Dr. Jean Davignon had observed a strikingly high

frequency of chylomicronemia. In collaboration with Dr. Hayden's team, Dr. Davignon and collaborators in Quebec, including Drs. C. Gagne and J. Bergeron, indeed found that the prevalence of LPL deficiency in regions of Quebec was very high (1 in 6,382 affected) compared to the rest of the world (<1 in 100,000) (Bergeron et al., 1992; Dionne et al., 1992; Wood et al., 1993; Pimstone et al., 1995; Marcil et al., 1996; Bijvoet et al., 1996a; Foubert et al., 1997b). Genealogical reconstruction revealed a genetic founder effect from early settlers to Quebec in the seventeenth century from France and Scotland had resulted in the highest worldwide frequency of LPL deficiency patients (Normand et al., 1992). Mutation analyses revealed two predominant LPL gene mutations in French-Canadian patients, G188E (20-30%) and P207L (~70%). Importantly, the common LPL gene mutations in Quebec were found to inactivate enzyme activity, leaving inactive LPL protein in the circulation (Monsalve et al., 1990; Normand et al., 1992). Dr. Hayden recognized that this might be important for therapeutic purposes. If one could deliver a new supply of functional LPL protein into patients, a "foreign" protein (i.e., normally absent in patients) would likely be detected by the patient's immune system and rapidly cleared. In contrast, in patients with detectable levels of inactive LPL protein, a newly supply of small amounts of therapeutic LPL enzyme administered to the patient could potentially function normally to overcome the enzyme deficit and reverse the disease symptoms.

Together with investigators around the world, novel LPL assays and antibodies to LPL were developed to improve the understanding of LPL biochemistry and function. Dr. Hayden's team regularly met with Dr. John Brunzell's Seattle-based research group, who developed critical antibodies to quantify LPL protein levels. Dr. Thomas Olivecrona's laboratory in Sweden significantly improved the understanding of the LPL enzyme. LPL enzyme is normally expressed in adipose tissue, skeletal muscle and heart muscle where it is secreted and attaches to the luminal surface of endothelial cells in capillaries. LPL breaks down triglycerides packaged in chylomicrons and very-low density lipoproteins to generate free fatty acids, which serve as a source of energy in muscle cells, or stored in adipose tissue (Levy et al., 1990; Liu and Olivecrona, 1992; Liu et al., 1993; Goldberg and Merkel, 2001)

Dr. Kastelein returned to the Netherlands to found the Lipid Research Clinic at the Academic Medical Centre in Amsterdam. The collaboration between Drs. Kastelein and Hayden continued with an ongoing identification of LPL deficient patients in Canada and the Netherlands and ongoing research into the function of LPL.

The research in Dr. Hayden's laboratory then began to focus on therapeutic strategies for LPL deficiency. A mouse model of LPL deficiency was developed by Dr. Clay Semenkovich in Washington University, St. Louis, and kindly provided to Dr. Hayden. One therapeutic strategy was to produce and purify large amounts of LPL enzyme and deliver this back into patients by frequent

intravenous injections. However, pre-clinical studies of intravenous LPL enzyme replacement therapy proved unfeasible due to the short *in vivo* half-life of the LPL protein (<10 min) (Liu et al., 2000).

Dr. Hayden's team began to explore gene therapy for LPL deficiency. The goal of gene therapy, first conceptualized as early as 1947, is to correct a human disease through the delivery of a functional therapeutic gene into patients (Keeler, 1947). In 1968, the use of viruses modified to carry genetic information were first developed in plants, and by 1972, Dr. Theodore Friedmann and Dr. Richard Roblin had predicted that gene therapy could potentially treat human diseases in the future (Friedmann and Roblin, 1972). In 1990, the first gene therapy proof-of-principle was achieved with the successful treatment of a rare immunodeficiency disease (adenosine deaminase deficiency) using a retroviral vector to deliver the therapeutic adenosine deaminase gene (Blaese et al., 1995). Building upon advances in gene therapy vectors, Dr. Hayden's team, including Dr. Suzanne Lewis, Dr. Kate Excoffon, and Dr. Guoqing Liu investigated a number of modified retroviral and adenoviral vectors to deliver functional copies of the LPL gene into cells (Lewis et al., 1995; Liu et al., 1997) and animal models (Excoffon et al., 1997). However, these preliminary studies were unable to achieve a sufficient duration or level of therapeutic LPL gene expression necessary to benefit patients.

A post-doctoral research fellow, Dr. Colin Ross, joined Dr. Hayden's laboratory to investigate the adeno-associated viral (AAV) vector. Drs. Ross and Hayden selected AAV because this was a non-pathogenic virus, incapable of replication, and carried no viral genes that could potentially trigger an immune-mediated clearance of treated cells. In essence, the modified AAV vector was a protein shell designed to efficiently and safely deliver the therapeutic LPL gene into the muscle cells of patients, where the LPL gene would generate an ongoing supply of therapeutic LPL enzyme (Ross et al., 2004). The initial results of AAV-mediated LPL gene delivery into cell culture were promising; however, the first several attempts to treat LPL deficient mice did not initially correct the hypertriglyceridemia.

A series of modifications were made in efforts to improve the effectiveness of the LPL gene therapy vector. Instead of using the wild-type (common) version of the LPL gene, a naturally occurring variant (S447X) was investigated. The S447X version of the LPL gene, present in 10-25% of people, had been associated with a significantly improved lipid profile and enhanced removal of lipoprotein particles from the circulation (Hata et al., 1990; Stocks et al., 1992; Jemaa et al., 1995; Galton et al., 1996; Gagne et al., 1999; Wittrup et al., 1999; 2002). Dr. Kastelein's team in the Netherlands, including Dr. Jan Albert Kuivenhoven, confirmed the improved lipid profile and enhanced plasma triglyceride reductions in people that carried the S447X variant (Groenemeijer et al., 1997a; Kuivenhoven et al., 1997; Groenemeijer et al., 1997b; Gagne et al., 1999; Henderson et al., 1999; Nierman et al., 2005). Back in Dr. Hayden's lab, a direct

comparison of the wild-type and S447X versions of the LPL gene's ability to rescue LPL deficient mice revealed that the S447X variant would be considerably more effective for gene therapy (0% rescue with wild-type versus 95% rescue with S447X) (Ross et al., 2005). The AAV capsid shell was also modified to alter the vector's tropism based upon reports from Dr. James Wilson that AAV serotype 1, although much less effective in cell culture, was significantly more effective in muscle than previous vectors (Gao et al., 2002).

After a series of vector modifications, on January 10, 2002, Dr. Ross administered the newest formulation (AAV1-LPL<sup>S447X</sup>) to the muscles of LPL deficient mice. Two weeks later, while drawing the blood samples, it was immediately apparent that the normally extremely high triglyceride levels of the LPL deficient mice had been significantly reduced based upon the appearance of the blood. As the blood samples were analyzed in the lab, the results became clear – the appearance of the plasma had changed from a pre-treatment opaque, milky-white to a transparent, normal-looking, plasma as featured on the cover of *Human Gene Therapy* (Fig. 3) (Ross et al., 2004). The blood triglyceride levels had been reduced 25-fold to near-normal levels. The LPL enzyme was being synthesized within the muscle and entering the circulation. Unlike previous attempts to deliver the LPL gene into mice, this time the treatment showed long-lasting effects (Ross et al., 2004). Blood samples collected month after month continued to show a near complete normalization of plasma triglyceride levels (Ross et al., 2004). Dr. Hayden delivered the good news to Dr. Kastelein in person in the Netherlands.

Drs. Hayden and Kastelein recognized that AAV1-LPL<sup>S447X</sup> could be a viable therapeutic option for LPL deficiency if the effects could be replicated in a large animal model and LPL deficient patients. Dr. Ross teamed up with Dr. Jaap Twisk of Amsterdam Molecular Therapeutics (AMT), a company that Dr. Kastelein had helped establish in the Netherlands, to optimize the treatment in mice and produce sufficient quantities of clinical-grade AAV for large animal studies and, optimistically, for clinical trials.

For many diseases, large animal models are not available. However, for LPL deficiency, there was a large animal model that Dr. Hayden had helped identify. Dr. Hayden had travelled to the Lipid and Drug Conference in Florence, Italy. During a conference poster session, Dr. Hayden met Dr. Boyd Jones, a veterinarian-researcher from Palmerston North, New Zealand, who was presenting a poster about the lipid anomalies in a domestic cat. Dr. Hayden noticed a remarkable similarity between the lipid anomalies of Dr. Boyd's cats and those of his LPL deficient patients. Working together, Drs. Hayden and Jones confirmed the LPL deficiency in the cats and identified the causal mutation in the LPL gene that resulted in a phenotype very similar to human LPL deficiency (Peritz et al., 1990; Ginzinger et al., 1996).

The next step was to determine whether the treatment would be effective in the larger cat model of LPL deficiency. With outstanding support from the University veterinarians and animal care staff, including Drs. Godbey, Love, and Harvey-Clark and G. Gray, Drs. Ross and Twisk administered a low dose of AAV1-LPL<sup>S447X</sup> to LPL deficient cats. Disappointingly, the initial results showed no effect. Using the remainder of a limited supply of the AAV vector, a nearly 20-fold higher dose was administered to the cats and the effects became immediately clear. The plasma triglyceride levels were normalized within a week and long-term correction was achieved (Ross et al., 2006). In LPL deficient cats, long-term correction required co-administration of an immunosuppressant due to the significant human-feline species differences in the LPL protein (Ross et al., 2006).

With proof-of-principle demonstrated in both small and large animal models, the next steps towards clinical application were large *in vivo* toxicological and biodistribution studies in animal models. These extensive studies demonstrated that the intramuscular administration of the AAV1-LPL<sup>S447X</sup> vector was well tolerated in animal models (Rip et al., 2005). In addition, clinical feasibility was demonstrated in cells harvested from muscle biopsies of LPL deficient patients. The patient's muscle cells were efficiently transduced by AAV1-LPL<sup>S447X</sup> and secreted active LPL (Rip et al., 2005). Together, these data supported the regulatory approval for a clinical trial of AAV1-LPL<sup>S447X</sup> in LPL-deficient patients (Rip et al., 2005), and Drs. Ross and Twisk confirmed the biological activity of the vector prepared for the clinical trial.

AMT supported the first clinical trial in the Netherlands led by Dr. Erik Stroes with patients that were originally identified by the Amsterdam and Vancouver teams. All of the patients had genetically-confirmed LPL deficiency with detectable LPL protein in their circulation. Highlighting the severity of LPL deficiency, during the pre-treatment monitoring phase of the trial, two of the patients developed severe pancreatitis (Rip et al., 2005). One patient recovered after receiving intensive care treatment, while the second, a 25-year-old patient, died as a direct result of the pancreatitis (Rip et al., 2005).

The first LPL deficient patient was treated with LPL gene therapy on Oct. 3, 2005. Under ultrasound guidance, patients were treated with either  $1 \times 10^{11}$  (n=4) or  $3 \times 10^{11}$  (n=4) genome copies (gc) per kg body weight via 40 or 60 intramuscular injections, respectively (Stroes et al., 2008). After 12 weeks, all patients had demonstrated a significant decrease of median triglyceride levels compared to baseline (27% and 41% reductions, respectively) (Stroes et al., 2008). No serious adverse events occurred and no antibody responses to LPL were observed (Stroes et al., 2008). The results also revealed that an immunosuppressive compound should be co-administered with the vector to minimize an immune response to AAV capsid proteins, and that a higher dose should be administered in future clinical trials (Stroes et al., 2008).

The team recognized that the requirement for higher doses posed a significant problem. Up until this time, the AAV had been produced with a plasmid-based system using adherent human embryonic kidney cells. However, the quantity of AAV produced in this system was insufficient for the requirements of a larger clinical trial at higher doses. Together with Dr. Wim Hermens, a scalable baculovirus-based AAV production system that used insect cells grown at high density in suspension culture was optimized to produce high quantities of AAV1-LPL<sup>S447X</sup>, now called Glybera® (alipogene tiparvovec). Drs. Ross and Twisk confirmed the therapeutic activity of the new baculovirus-generated AAV formulation and a new, larger, clinical trial received regulatory approval to proceed with higher doses.

The story of the development of Glybera® returned once again to the province of Quebec, Canada, in recognition that Quebec had the highest worldwide frequency of LPL deficiency in the world, and building upon the collaborations that had been initiated years earlier (Pimstone et al., 1995). The second LPL gene therapy clinical trial was conducted in Quebec, led by Dr. Daniel Gaudet at the Université de Montréal. LPL deficient patients with a history of pancreatitis were treated with  $3 \times 10^{11}$  gc/kg or  $1 \times 10^{12}$  gc/kg of AAV1-LPL<sup>S447X</sup> administered in multiple intramuscular injections into leg muscles, as well as a 12 week regimen of immunosuppressants cyclosporine A (3mg/kg per day) and mycophenolate mofetil (2 g per day). The trial also included a bridging sub-study that demonstrated similar effects of AAV1-LPL<sup>S447X</sup> generated using the plasmid system versus the baculovirus system. Overall, the treatment was well tolerated, without emerging safety concerns for over two years following treatment (Gaudet et al., 2012b). By 12 weeks, plasma triglycerides had been significantly reduced an average 40% with reductions of up to 77% observed. After six months, sustained LPL<sup>S447X</sup> expression and long-term changes in triglyceride-rich lipoprotein characteristics were noted, independent of effects on plasma triglycerides (Gaudet et al., 2012b). After two years, more than half (64%) of the patients reported clinical benefits, including a capacity to eat more, or to eat food they were previously unable to eat, and increased energy levels or improved abdominal comfort (Gaudet et al., 2012b). Importantly, the incidence of pancreatitis was significantly reduced five-fold.

A small follow-up clinical study was conducted in Quebec to document the postprandial effects of Glybera®, and collect additional efficacy and safety data using the same dose and immunosuppression regimen as before ( $1 \times 10^{12}$  gc/kg) (Carpentier et al., 2012). At 14 weeks post-treatment, intramuscular administration of Glybera® significantly improved postprandial chylomicron metabolism by 79 and 93%, 6h and 24h after a test meal, respectively. One of the participants in this study became pregnant a year post-treatment. Pregnancy is an important trigger of pancreatitis and can be life-threatening in LPL deficiency (Ma et al., 1993; Ewald et al., 2009); however, for this treated patient the pregnancy went very well and she delivered a healthy baby.

A blinded retrospective and prospective review of treated and untreated LPL deficient patients was recently conducted to generate an unbiased assessment of the incidence of pancreatitis before and after Glybera® treatment. The results thus far, have shown reductions in the incidence, duration, and severity of pancreatitis episodes, as well as a reduction in the number and duration of hospitalizations (Gaudet et al., 2012a).

After a series of significant delays, the European Medicines Agency granted regulatory approval for Glybera® on November 2, 2012, making it the first gene therapy to receive regulatory approval in the Western World (Moran, 2012).

The story of Glybera® involved the dedicated contributions of many patients, students, fellows, lab technologists, clinicians, and researchers from around the world. The journey consisted of many lifetimes of work, which led to the creation of many lifelong friendships and ongoing research collaborations. Building upon the significant pre-clinical research, the development path for Glybera® was relatively fast. The application for regulatory approval of Glybera® was submitted December 23, 2009, four years after the first administration in humans. However, the regulatory approval of Glybera® was excruciatingly slow. Nevertheless, it is hoped that the approval of this novel gene therapy will help open the doors to new gene therapies that will benefit patients in the future.

### **Acknowledgments**

We would like to acknowledge the participation of all of the patients and their families involved at all stages of the process of the development of LPL gene therapy. We also want to acknowledge the support of all of the students and postdoctoral fellows who contributed over many years towards the development of LPL gene therapy, and the staff of AMT who all worked so diligently to make this happen. We also want to acknowledge the support of our families and friends throughout this journey.

### **Author Disclosure Statement**

JJPK: consultant to UniQure.

CJDR: No competing financial interests exist.


**FIGURE LEGEND:**



**Figure 1. Cutaneous xanthomas of Lipoprotein Lipase Deficiency. (Image courtesy of Dr. Kastelein).**



**Figure 2. Michael Hayden (third from left) and John Kastelein (far right) and the Hayden Lab team toasting on the identification of the first mutation in the LPL gene.**



**Figure 3. Cover courtesy of *Human Gene Therapy* showing the effect of AAV1-LPL<sup>S447X</sup> on plasma lipidemia over time. (Ross et al., 2004).**

## REFERENCES

- Bergeron, J., Normand, T., Bharucha, A., Ven Murthy, M.R., Julien, P., Gagne, C., Dionne, C., De Braekeleer, M., Brun, D., and Hayden, M.R. (1992). Prevalence, geographical distribution and genealogical investigations of mutation 188 of lipoprotein lipase gene in the French Canadian population of Quebec. *Clin. Genet.* 41, 206–210.
- Bijvoet, S., Gagne, S.E., Moorjani, S., Gagne, C., Henderson, H.E., Fruchart, J.C., Dallongeville, J., Alaupovic, P., Prins, M., Kastelein, J.J., et al. (1996a). Alterations in plasma lipoproteins and apolipoproteins before the age of 40 in heterozygotes for lipoprotein lipase deficiency. *J. Lipid Res.* 37, 640–650.
- Bijvoet, S.M., Bruin, T., Tuzgöl, S., Bakker, H.D., Hayden, M.R., and Kastelein, J.J. (1994). Homozygosity for a mutation in the lipoprotein lipase gene (Gly139-->Ser) causes chylomicronaemia in a boy of Spanish descent. *Hum. Genet.* 93, 339–343.
- Bijvoet, S.M., Wiebusch, H., Ma, Y., Reymer, P.W., Bruin, T., Bakker, H.D., Funke, H., Assmann, G., Hayden, M.R., and Kastelein, J.J. (1996b). Compound heterozygosity for a known and a novel defect in the lipoprotein lipase gene (Asp250-->Asn; Ser251-->Cys) resulting in lipoprotein lipase (LPL) deficiency. *Neth J Med* 49, 189–195.
- Blaese, R.M., Culver, K.W., Miller, A.D., Carter, C.S., Fleisher, T., Clerici, M., Shearer, G., Chang, L., Chiang, Y., Tolstoshev, P., et al. (1995). T lymphocyte-directed gene therapy for ADA- SCID: initial trial results after 4 years. *Science* (New York, N.Y) 270, 475–480.
- Bruin, T., Kastelein, J.J., van Diermen, D.E., Ma, Y., Henderson, H.E., Stuyt, P.M., Stalenhoef, A.F., Sturk, A., Brunzell, J.D., and Hayden, M.R. (1992). A missense mutation Pro157 Arg in lipoprotein lipase (LPLNijmegen) resulting in loss of catalytic activity. *Eur. J. Biochem.* 208, 267–272.
- Bruin, T., Tuzgöl, S., Mulder, W.J., van den Ende, A.E., Jansen, H., Hayden, M.R., and Kastelein, J.J. (1994). A compound heterozygote for lipoprotein lipase deficiency, Val69-->Leu and Gly188-->Glu: correlation between in vitro LPL activity and clinical expression. *J. Lipid Res.* 35, 438–445.
- Brunzell, J., Deeb, S., Scriver, C., Beaudet, A., Sly, W., and Valle, D. (2001). Familial lipoprotein lipase deficiency, ApoCII deficiency, and hepatic lipase deficiency. In *The Metabolic and Molecular Bases of Inherited Disease*, (New York, NY: McGraw-Hill Inc), pp. 2789–2816.
- Carpentier, A.C., Frisch, F., Labb  , S.M., Gagnon, R., de Wal, J., Greentree, S., Petry, H., Twisk, J., Brisson, D., and Gaudet, D. (2012). Effect of alipogene tiparvovec (AAV1-LPL(S447X)) on postprandial chylomicron metabolism in lipoprotein lipase-deficient patients. *J Clin Endocrinol Metab* 97, 1635–1644.

- Devlin, R.H., Deeb, S., Brunzell, J., and Hayden, M.R. (1990). Partial gene duplication involving exon-Alu interchange results in lipoprotein lipase deficiency. *American Journal of Human Genetics* 46, 112–119.
- Dionne, C., Gagne, C., Julien, P., Murthy, M.R., Lambert, M., Roederer, G., Davignon, J., Hayden, M.R., Lupien, P.J., and De Braekeleer, M. (1992). Genetic epidemiology of lipoprotein lipase deficiency in Saguenay-Lac-St-Jean (Québec, Canada). *Ann. Genet.* 35, 89–92.
- Ewald, N., Hardt, P.D., and Kloer, H.-U. (2009). Severe hypertriglyceridemia and pancreatitis: presentation and management. *Curr. Opin. Lipidol.* 20, 497–504.
- Excoffon, K.J., Liu, G., Miao, L., Wilson, J.E., McManus, B.M., Semenkovich, C.F., Coleman, T., Benoit, P., Duverger, N., Branellec, D., et al. (1997). Correction of hypertriglyceridemia and impaired fat tolerance in lipoprotein lipase-deficient mice by adenovirus-mediated expression of human lipoprotein lipase. *Arterioscler. Thromb. Vasc. Biol.* 17, 2532–2539.
- Foubert, L., Bruin, T., De Gennes, J.L., Ehrenborg, E., Furioli, J., Kastelein, J., Benlian, P., and Hayden, M. (1997a). A single Ser259Arg mutation in the gene for lipoprotein lipase causes chylomicronemia in Moroccans of Berber ancestry. *Hum Mutat* 10, 179–185.
- Foubert, L., De Gennes, J.L., Lagarde, J.P., Ehrenborg, E., Raisonniere, A., Girardet, J.P., Hayden, M.R., and Benlian, P. (1997b). Assessment of French patients with LPL deficiency for French Canadian mutations. *J. Med. Genet.* 34, 672–675.
- Friedmann, T., and Roblin, R. (1972). Gene therapy for human genetic disease? *Science* (New York, N.Y) 175, 949–955.
- Gagne, S.E., Larson, M.G., Pimstone, S.N., Schaefer, E.J., Kastelein, J.J., Wilson, P.W., Ordovas, J.M., and Hayden, M.R. (1999). A common truncation variant of lipoprotein lipase (Ser447X) confers protection against coronary heart disease: the Framingham Offspring Study. *Clin. Genet.* 55, 450–454.
- Galton, D.J., Mattu, R., Needham, E.W., and Cavanna, J. (1996). Identification of putative beneficial mutations for lipid transport. *Z. Gastroenterol.* 34 Suppl 3:56–8, 56–58.
- Gao, G.P., Alvira, M.R., Wang, L., Calcedo, R., Johnston, J., and Wilson, J.M. (2002). Novel adeno-associated viruses from rhesus monkeys as vectors for human gene therapy. *Proceedings of the National Academy of Sciences of the United States of America* 99, 11854–11859.
- Gaudet, D., Méthot, J., and Kastelein, J. (2012a). Gene therapy for lipoprotein lipase deficiency. *Curr. Opin. Lipidol.* 23, 310–320.

- Gaudet, D., Méthot, J., Déry, S., Brisson, D., ESSIEMBRE, C., Tremblay, G., Tremblay, K., de Wal, J., Twisk, J., van den Bulk, N., et al. (2012b). Efficacy and long-term safety of alipogene tiparvovec (AAV1-LPL(S447X)) gene therapy for lipoprotein lipase deficiency: an open-label trial. *Gene Ther.*
- Ginzinger, D.G., Lewis, M.E., Ma, Y., Jones, B.R., Liu, G., and Jones, S.D. (1996). A mutation in the lipoprotein lipase gene is the molecular basis of chylomicronemia in a colony of domestic cats. *The Journal of Clinical Investigation* 97, 1257–1266.
- Goldberg, I.J., and Merkel, M. (2001). Lipoprotein lipase: physiology, biochemistry, and molecular biology. *Front. Biosci.* 6, D388–D405.
- Groenemeijer, B.E., Hallman, M.D., Reymer, P.W., Gagne, E., Kuivenhoven, J.A., Bruin, T., Jansen, H., Lie, K.I., Bruschke, A.V., Boerwinkle, E., et al. (1997a). Genetic variant showing a positive interaction with beta-blocking agents with a beneficial influence on lipoprotein lipase activity, HDL cholesterol, and triglyceride levels in coronary artery disease patients. The Ser447-stop substitution in the lipoprotein lipase gene. REGRESS Study Group. *Circulation* 95, 2628–2635.
- Groenemeijer, B.E., Hallman, M.D., Reymer, P.W., Gagne, E., Kuivenhoven, J.A., Bruin, T., Jansen, H., Lie, K.I., Bruschke, A.V., Boerwinkle, E., et al. (1997b). Genetic variant with a beneficial influence on lipoprotein lipase activity, HDL cholesterol, and triglyceride levels in coronary artery disease patients. *Circulation* 95, 2628–2635.
- Hata, A., Robertson, M., Emi, M., and Lalouel, J.M. (1990). Direct detection and automated sequencing of individual alleles after electrophoretic strand separation: identification of a common nonsense mutation in exon 9 of the human lipoprotein lipase gene. *Nucleic Acids Research* 18, 5407–5411.
- Henderson, H., Leisegang, F., Hassan, F., Hayden, M., and Marais, D. (1998a). A novel Glu421Lys substitution in the lipoprotein lipase gene in pregnancy-induced hypertriglyceridemic pancreatitis. *Clinica Chimica Acta; International Journal of Clinical Chemistry* 269, 1–12.
- Henderson, H.E., Bijvoet, S.M., Mannens, M.A., Bruin, T., Erkelens, D.W., Hayden, M.R., and Kastelein, J.J. (1998b). Ile225Thr loop mutation in the lipoprotein lipase (LPL) gene is a de novo event. *Am. J. Med. Genet.* 78, 313–316.
- Henderson, H.E., Devlin, R., Peterson, J., Brunzell, J.D., and Hayden, M.R. (1990). Frameshift mutation in exon 3 of the lipoprotein lipase gene causes a premature stop codon and lipoprotein lipase deficiency. *Mol. Biol. Med.* 7, 511–517.
- Henderson, H.E., Hassan, F., Berger, G.M., and Hayden, M.R. (1992). The

lipoprotein lipase Gly188----Glu mutation in South Africans of Indian descent: evidence suggesting common origins and an increased frequency. *J. Med. Genet.* 29, 119–122.

Henderson, H.E., Kastelein, J.J., Zwinderman, A.H., Gagne, E., Jukema, J.W., Reymer, P.W., Groenemeyer, B.E., Lie, K.I., Bruschke, A.V., Hayden, M.R., et al. (1999). Lipoprotein lipase activity is decreased in a large cohort of patients with coronary artery disease and is associated with changes in lipids and lipoproteins. *J. Lipid Res.* 40, 735–743.

Jemaa, R., Fumeron, F., Poirier, O., Lecerf, L., Evans, A., Arveiler, D., Luc, G., Cambou, J.P., Bard, J.M., and Fruchart, J.C. (1995). Lipoprotein lipase gene polymorphisms: associations with myocardial infarction and lipoprotein levels, the ECTIM study. *Etude Cas Temoin sur l'Infarctus du Myocarde. J. Lipid Res.* 36, 2141–2146.

Jukema, J.W., van Boven, A.J., Groenemeijer, B., Zwinderman, A.H., Reiber, J.H., Bruschke, A.V., Henneman, J.A., Molhoek, G.P., Bruin, T., Jansen, H., et al. (1996). The Asp9 Asn mutation in the lipoprotein lipase gene is associated with increased progression of coronary atherosclerosis. REGRESS Study Group, Interuniversity Cardiology Institute, Utrecht, The Netherlands. *Regression Growth Evaluation Statin Study. Circulation* 94, 1913–1918.

Kastelein, J.J., Groenemeyer, B.E., Hallman, D.M., Henderson, H., Reymer, P.W., Gagne, S.E., Jansen, H., Seidell, J.C., Kromhout, D., Jukema, J.W., et al. (1998). The Asn9 variant of lipoprotein lipase is associated with the -93G promoter mutation and an increased risk of coronary artery disease. The Regress Study Group. *Clin. Genet.* 53, 27–33.

Keeler, C.E. (1947). Gene therapy. *J. Hered.* 38, 294–298.

Kuivenhoven, J.A., Groenemeyer, B.E., Boer, J.M., Reymer, P.W., Berghuis, R., Bruin, T., Jansen, H., Seidell, J.C., and Kastelein, J.J. (1997). Ser447stop mutation in lipoprotein lipase is associated with elevated HDL cholesterol levels in normolipidemic males. *Arterioscler. Thromb. Vasc. Biol.* 17, 595–599.

Langlois, S., Deeb, S., Brunzell, J.D., Kastelein, J.J., and Hayden, M.R. (1989). A major insertion accounts for a significant proportion of mutations underlying human lipoprotein lipase deficiency. *Proceedings of the National Academy of Sciences of the United States of America* 86, 948–952.

Levy, E., Deckelbaum, R.J., Thibault, R.L., Seidman, E., Olivecrona, T., and Roy, C.C. (1990). In vitro remodelling of plasma lipoproteins in whole plasma by lipoprotein lipase in primary and secondary hypertriglyceridaemia. *Eur. J. Clin. Invest.* 20, 422–431.

Lewis, M.E., Forsythe, I.J., Marth, J.D., Brunzell, J.D., Hayden, M.R., and Humphries, R.K. (1995). Retroviral-mediated gene transfer and expression of

- human lipoprotein lipase in somatic cells. *Hum. Gene Ther.* 6, 853–863.
- Liu, G., and Olivecrona, T. (1992). Synthesis and transport of lipoprotein lipase in perfused guinea pig hearts. *Am. J. Physiol.* 263, H438–H446.
- Liu, G., Ashbourne, E.K.J., Wilson, J.E., McManus, B.M., Rogers, Q.R., Miao, L., Kastelein, J.J., Lewis, M.E., and Hayden, M.R. (2000). Phenotypic correction of feline lipoprotein lipase deficiency by adenoviral gene transfer. *Hum. Gene Ther.* 11, 21–32.
- Liu, G., Bengtsson-Olivecrona, G., and Olivecrona, T. (1993). Assembly of lipoprotein lipase in perfused guinea-pig hearts. *Biochem. J.* 292 (Pt 1), 277–282.
- Liu, G., Excoffon, K.J., Benoit, P., Ginzinger, D.G., Miao, L., Ehrenborg, E., Duverger, N., Denefle, P.P., Hayden, M.R., and Lewis, M.E. (1997). Efficient adenovirus-mediated ectopic gene expression of human lipoprotein lipase in human hepatic (HepG2) cells. *Hum. Gene Ther.* 8, 205–214.
- Ma, Y., Liu, M.S., Ginzinger, D., Frohlich, J., Brunzell, J.D., and Hayden, M.R. (1993). Gene-environment interaction in the conversion of a mild-to-severe phenotype in a patient homozygous for a Ser172→Cys mutation in the lipoprotein lipase gene. *The Journal of Clinical Investigation* 91, 1953–1958.
- Ma, Y.H., Bruin, T., Tuzgöl, S., Wilson, B.I., Roederer, G., Liu, M.S., Davignon, J., Kastelein, J.J., Brunzell, J.D., and Hayden, M.R. (1992). Two naturally occurring mutations at the first and second bases of codon aspartic acid 156 in the proposed catalytic triad of human lipoprotein lipase. In vivo evidence that aspartic acid 156 is essential for catalysis. *The Journal of Biological Chemistry* 267, 1918–1923.
- Marcil, M., Boucher, B., Gagne, E., Davignon, J., Hayden, M., and Genest, J. (1996). Lack of association of the apolipoprotein A-I-C-III-A-IV gene Xmnl and SstI polymorphisms and of the lipoprotein lipase gene mutations in familial combined hyperlipoproteinemia in French Canadian subjects. *J. Lipid Res.* 37, 309–319.
- Monsalve, M.V., Henderson, H., Roederer, G., Julien, P., Deeb, S., Kastelein, J.J., Peritz, L., Devlin, R., Bruin, T., and Murthy, M.R. (1990). A missense mutation at codon 188 of the human lipoprotein lipase gene is a frequent cause of lipoprotein lipase deficiency in persons of different ancestries. *The Journal of Clinical Investigation* 86, 728–734.
- Moran, N. (2012). First gene therapy approved. *Nat. Biotechnol.* 30, 1153.
- Nierman, M.C., Prinsen, B.H., Rip, J., Veldman, R.J., Kuivenhoven, J.A., Kastelein, J.J., de Sain-van der Velden, M.G., and Stroes, E.S. (2005). Enhanced Conversion of Triglyceride-Rich Lipoproteins and Increased Low-

Density Lipoprotein Removal in LPLS447X Carriers. *Arterioscler. Thromb. Vasc. Biol.* .

- Normand, T., Bergeron, J., Fernandez-Margallo, T., Bharucha, A., Ven Murthy, M.R., Julien, P., Gagne, C., Dionne, C., De Braekeleer, M., and Ma, R. (1992). Geographic distribution and genealogy of mutation 207 of the lipoprotein lipase gene in the French Canadian population of Quebec. *Hum. Genet.* 89, 671–675.
- Peritz, L.N., Brunzell, J.D., Harvey-Clarke, C., Pritchard, P.H., Jones, B.R., and Hayden, M.R. (1990). Characterization of a lipoprotein lipase class III type defect in hypertriglyceridemic cats. *Clin Invest Med* 13, 259–263.
- Pimstone, S.N., Clee, S.M., Gagne, S.E., Miao, L., Zhang, H., Stein, E.A., and Hayden, M.R. (1996). A frequently occurring mutation in the lipoprotein lipase gene (Asn291Ser) results in altered postprandial chylomicron triglyceride and retinyl palmitate response in normolipidemic carriers. *J. Lipid Res.* 37, 1675–1684.
- Pimstone, S.N., Gagne, S.E., Gagne, C., Lupien, P.J., Gaudet, D., Williams, R.R., Kotze, M., Reymer, P.W., Defesche, J.C., and Kastelein, J.J. (1995). Mutations in the gene for lipoprotein lipase. A cause for low HDL cholesterol levels in individuals heterozygous for familial hypercholesterolemia. *Arterioscler. Thromb. Vasc. Biol.* 15, 1704–1712.
- Rip, J., Nierman, M.C., Sierts, J.A., Petersen, W., Van Den Oever, K., Van Raalte, D., Ross, C.J., Hayden, M.R., Bakker, A.C., Dijkhuizen, P., et al. (2005). Gene therapy for lipoprotein lipase deficiency: working toward clinical application. *Hum. Gene Ther.* 16, 1276–1286.
- Rip, J., Nierman, M.C., Sierts, J.A., Van Raalte, D., Petersen, W., Van Den Oever, K., Ross, C.J., Hayden, M.R., Meulenbergh, J.M., Twisk, J., et al. (2003). Working towards clinical application of gene therapy for LPL deficiency. *Circulation* 108, 133–133.
- Ross, C.J.D., Liu, G.Q., Kuivenhoven, J.A., Twisk, J., Rip, J., van Dop, W., Excoffon, K., Lewis, S.M.E., Kastelein, J.J., and Hayden, M.R. (2005). Complete rescue of lipoprotein lipase-deficient mice by somatic gene transfer of the naturally occurring LPLS447X beneficial mutation. *Arterioscler. Thromb. Vasc. Biol.* 25, 2143–2150.
- Ross, C.J.D., Twisk, J., Bakker, A.C., Miao, F., Verbart, D., Rip, J., Godbey, T., Dijkhuizen, P., Hermens, W., Kastelein, J.J.P., et al. (2006). Correction of feline lipoprotein lipase deficiency with adeno-associated virus serotype 1-mediated gene transfer of the lipoprotein lipase S447X beneficial mutation. *Hum. Gene Ther.* 17, 487–499.
- Ross, C.J.D., Twisk, J., Meulenbergh, J.M., Liu, G., Van den Oever, K., Moraal, E., Hermens, W.T., Rip, J., Kastelein, J.J.P., Kuivenhoven, J.A., et al. (2004).

Long-term correction of murine lipoprotein lipase deficiency with AAV1-mediated gene transfer of the naturally occurring LPL(S447X) beneficial mutation. *Hum. Gene Ther.* 15, 906–919.

Stocks, J., Thorn, J.A., and Galton, D.J. (1992). Lipoprotein lipase genotypes for a common premature termination codon mutation detected by PCR-mediated site-directed mutagenesis and restriction digestion. *J. Lipid Res.* 33, 853–857.

Stroes, E.S., Nierman, M.C., Meulenbergh, J.J., Franssen, R., Twisk, J., Henny, C.P., Maas, M.M., Zwinderman, A.H., Ross, C., Aronica, E., et al. (2008). Intramuscular administration of AAV1-lipoprotein lipase S447X lowers triglycerides in lipoprotein lipase-deficient patients. *Arterioscler. Thromb. Vasc. Biol.* 28, 2303–2304.

Wion, K.L., Kirchgessner, T.G., Lusis, A.J., Schotz, M.C., and Lawn, R.M. (1987). Human lipoprotein lipase complementary DNA sequence. *Science* (New York, N.Y) 235, 1638–1641.

Wittekoek, M.E., Pimstone, S.N., Reymer, P.W., Feuth, L., Botma, G.J., Defesche, J.C., Prins, M., Hayden, M.R., and Kastelein, J.J. (1998). A common mutation in the lipoprotein lipase gene (N291S) alters the lipoprotein phenotype and risk for cardiovascular disease in patients with familial hypercholesterolemia. *Circulation* 97, 729–735.

Wittrup, H.H., Nordestgaard, B.G., Steffensen, R., Jensen, G., and Tybjaerg-Hansen, A. (2002). Effect of gender on phenotypic expression of the S447X mutation in LPL: the Copenhagen City Heart Study. *Atherosclerosis* 165, 119–126.

Wittrup, H.H., Tybjaerg-Hansen, A., and Nordestgaard, B.G. (1999). Lipoprotein lipase mutations, plasma lipids and lipoproteins, and risk of ischemic heart disease. A meta-analysis. *Circulation* 99, 2901–2907.

Wood, S., Schertzer, M., Hayden, M., and Ma, Y. (1993). Support for founder effect for two lipoprotein lipase (LPL) gene mutations in French Canadians by analysis of GT microsatellites flanking the LPL gene. *Hum. Genet.* 91, 312–316.